

Lyme Disease Incidence Reported by CDC – 2013*
(cases per 100,000 population)

Ranking	State		Incidence
1	VT	Vermont	107.6
2	NH	New Hampshire	100.0
3	ME	Maine	84.8
4	CT	Connecticut ▲	58.7
5	MA	Massachusetts	57.0
6	DE	Delaware	43.2
7	RI	Rhode Island ▲	42.2
8	PA	Pennsylvania	39.0
9	NJ	New Jersey	31.3
10	MN	Minnesota	26.4
11	WI	Wisconsin ▲	25.2
12	NY	New York ▲	17.9
13	MD	Maryland	13.5
14	VA	Virginia	11.2
15	WV	West Virginia	6.3
16	DC	Washington DC ▲	5.1
17	IA	Iowa	5.0
18	IL	Illinois	2.6
19	AK	Alaska	1.9
20	ND	North Dakota	1.7
21	MT	Montana	1.6
22	IN	Indiana	1.5
23	MI	Michigan	1.2
24	ID	Idaho	0.9
25	SC	South Carolina	0.7
26	KS	Kansas	0.6
26	OH	Ohio	0.6
27	FL	Florida	0.4
27	KY	Kentucky	0.4
27	NE	Nebraska	0.4
27	NV	Nevada	0.4
27	NC	North Carolina	0.4
27	SD	South Dakota	0.4
28	AZ	Arizona	0.3
28	OR	Oregon	0.3
28	UT	Utah	0.3
29	AL	Alabama	0.2
29	CA	California ▲	0.2
29	TN	Tennessee	0.2
29	TX	Texas	0.2
29	WA	Washington	0.2
29	WY	Wyoming	0.2
30	GA	Georgia	0.1
31	AK	Arkansas	0.0
31	CO	Colorado	0.0
31	LA	Louisiana	0.0
31	MS	Mississippi	0.0
31	MO	Missouri	0.0
31	NM	New Mexico	0.0
31	OK	Oklahoma	0.0
–	HI	Hawaii ▲	N
		U.S. Total	8.6

* CDC adopted a change in case definition in 2008. In addition to "Confirmed" cases, the "Probable" category was reported out for the first time. 2013 incidence includes only confirmed cases reported by CDC. US 2013 confirmed cases = 27,203 / probable = 9,104.

Note: CDC states only 10% of Lyme disease cases meeting the surveillance definition are reported – 10,000 cases are reported, 100,000 cases occurred (does not include all the cases falling outside the stringent surveillance case definition). In 2013, CDC announced that about 300,000 cases of Lyme actually occur annually.

Source: Data compiled from CDC published data (www.cdc.gov/lyme/stats/chartstables/incidencebystate.html)